

Pediatric turbinoplasty

Clinical Policy ID: CCP.1507

Recent review date: 3/2024

Next review date: 7/2025

Policy contains: nasal congestion, turbinate hypertrophy, rhinitis, rhinosinusitis, turbinectomy, turbinoplasty

AmeriHealth Caritas Next has developed clinical policies to assist with making coverage determinations. AmeriHealth Caritas Next's clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of "medically necessary," and the specific facts of the particular situation are considered by AmeriHealth Caritas Next when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. AmeriHealth Caritas Next's clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients. AmeriHealth Caritas Next's clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, AmeriHealth Caritas Next will update its clinical policies as necessary. AmeriHealth Caritas Next's clinical policies are not guarantees of payment.

Coverage policy

Turbinoplasty, or turbinate reduction, for children is clinically proven for persistent, symptomatic turbinate hypertrophy and, therefore, may be medically necessary when all of the following criteria are met:

- Turbinate hypertrophy is documented by direct visualization or nasal endoscopy.
- The enlarged turbinates are causing obstruction to breathing or chronic sinusitis (symptoms \geq 6 weeks).
- Intranasal corticosteroid spray has been trialed for at least 3 months and failed to adequately improve symptoms or is contraindicated/not tolerated (Maniaci, 2023).
- A course of medical management has been attempted and failed/medical management is contraindicated.
- Only inferior turbinates are reduced (National Institute for Health and Care Excellence, 2014; Seidman, 2015).

Turbinate reduction may be performed during tonsillectomy/adenoidectomy if the turbinates are enlarged and causing significant nasal obstruction.

Limitations

Children under age two are not candidates for turbinoplasty. The procedure is investigational/not clinically proven and therefore not medically necessary before this age.

Turbinectomy, or removal of bone, is not clinically proven and, therefore, not medically necessary.

Alternative covered services

- Balloon ostial sinuplasty.
- Standard treatments for chronic nasal congestion, deviated septum, snoring, sleep apnea, or nosebleeds, including corticosteroid injections and nasal corticosteroid sprays/decongestants.

Background

The superior, middle, and inferior turbinates are bonelike structures in the inferior part of the nose that clean, warm, and humidify inhaled air. Turbinate hypertrophy, which can be caused by allergy, infection, and hormonal changes, results in nasal obstruction (allergic, vasomotor, or infectious rhinitis), as measured by the amount of air flow in the nose (Abdullah, 2021). A study of 544 children with allergic rhinitis found 438 (80.8%) were diagnosed with turbinate hypertrophy (Ciprandi, 2020).

Antihistamines, topical decongestants, and topical corticosteroids are often effective in treating nasal obstruction. If these therapies are unsuccessful after three to six months, surgery can be considered (Komshian, 2019; Scheithauer, 2010). Procedures include adenoidectomy, sinus puncture/lavage, open surgical approaches, endoscopic sinus surgery, balloon sinuplasty, turbinectomy, and turbinate reduction (Isaacson, 2015).

Surgical reduction of the inferior turbinate can relieve nasal block while retaining turbinate function. Surgery of the inferior turbinate involves removal of the mucosa, soft erectile tissue, and turbinate bone. Techniques include conventional (partial or total), laser, cryo-, and electrocautery turbinectomy. Because surgery results in excess loss of bone and mucosa tissue, turbinoplasty (which preserves functional medial mucosa) can be performed instead (Abdullah, 2021), most commonly by submucous resection or tissue ablation (Seidman, 2015). The turbinates are composed of fleshy tissue with very fine bones supporting them so that the goal is not to remove bone but to decrease the fleshy tissue.

Turbinate reduction surgery in children, as compared with adults, has been described as “contentious and debatable” due to concerns over complications like excessive bleeding, damage to the mucosa with synechia and tear, disruption of nasal physiology and function, and disturbance of facial development (Abdullah, 2021).

One review states that “many surgeons feel it is prudent” to wait until the patient’s nasal growth is completed before surgery is performed (Gary, 2017). Another contends that “no ideal standard technique for turbinate reduction has been developed,” citing a lack of comparable long-term studies, but concedes anterior turbinoplasty to be the method of choice based on limited information (Scheithauer, 2010).

A review notes that turbinate reduction in the pediatric population can be achieved through turbinectomy, electrocautery, lasers, submucous microdebridement, and radiofrequency volumetric tissue reduction, but that consensus on the preferred approach is lacking (Komshian, 2019). In the pediatric population, the procedure involves reduction of excess reactive soft tissues of the turbinate and not the underlying supporting bone.

A recent review notes that while turbinate surgery in pediatric patients is gradually increasing in popularity amongst pediatric otolaryngologists... there is scarce information regarding this surgical procedure in children” (Calvo-Henriquez, 2022).

Findings

The American Association of Family Physicians’ guideline on rhinosinusitis only addresses adult cases (Sedaghat, 2017). The American Academy of Otolaryngology’s guideline on allergic rhinitis with nasal obstruction

and enlarged inferior turbinates supports the referral of patients who failed medical management to specialists for surgery; children younger than age 2 were excluded from this guideline. The Academy concludes that “inferior reduction surgery is a reasonable option” for allergic rhinitis patients who have inferior turbinate hypertrophy with symptoms despite medical management; benefits cited include improved symptoms, improved quality of life, improved medication delivery, reduced medication use, and better sleep. The Academy graded evidence for necessity Moderate/Grade C, due to a lack of head-to-head trials between medical and surgical outcomes (Seidman, 2015).

An American Academy of Pediatrics guideline on diagnosis and treatment of acute bacterial rhinosinusitis in children 1-18 years mentions that swelling of nasal turbinates are non-specific findings but does not mention turbinate surgery as an option (Wald, 2013).

A guideline from the United Kingdom states that, in cases of turbinate hypertrophy refractory to medical management, evidence supports the safety of radiofrequency tissue reduction for turbinate hypertrophy, and supports efficacy up to two years after surgery. However, the guideline includes microdebrider-assisted and laser-assisted turbinoplasty as acceptable, but not turbinectomy (National Institute for Health and Care Excellence, 2014).

A review recommended that pediatric inferior turbinate hypertrophy treatment should start with three months of medical management, evaluation for adenoid hypertrophy for consideration of concurrent adenoidectomy, and, finally, radiofrequency volumetric tissue reduction or submucous microdebridement as first-line surgery (Komshian, 2019).

In 2023, to address this gap and the absence of specific surgical guidelines for children, an international group of otolaryngologists developed a clinical consensus statement using a modified Delphi process, incorporating a literature review of eight articles including meta-analysis, systematic reviews, and case studies (n = 3, 088). The recommendations made pertained to the diagnosis, treatment, and follow-up of pediatric turbinate hypertrophy. They included guidance on using diagnostic tools like anterior rhinoscopy and fiberoptic examination, medical therapies like intranasal corticosteroids and saline irrigations, and surgical interventions such as pediatric turbinoplasty with minimally invasive techniques (Maniaci, 2023).

A literature review observed that non-mucosal-sparing turbinectomy surgery often results in postoperative complications (excessive bleeding, crusting, pain, and prolonged recovery period), and thus mucosal-sparing procedures (turbinoplasty) are typically the preferred option for cases of nasal obstruction refractory to conservative treatment (Abdullah, 2021).

A systematic review of 13 studies (n = 1,111) of turbinate surgery in children showed postoperative improvement in nasal congestion. Authors concluded the procedure to be safe, based on a complication rate of 3.12% — mostly minor bleeding, crust, and pain. Due to poor quality of studies, a surgical technique could not be recommended, but the safest are microdebrider-assisted inferior turbinoplasty, radiofrequency, coblation, and laser (Calvo-Henriquez, 2020).

A review of 1,770 children treated with an inferior turbinoplasty procedure at a Colorado hospital from 2003 to 2013 included 107 with an isolated procedure, namely radiofrequency ablation (72); microdebridement (19); and partial turbinate resection (21). None experienced major complications. Over half (54.0%) continued to require medical management; and authors note hypertrophy is not solely a surgical disease (Arganbright, 2015).

A systematic review of 58 studies analyzed results of surgery for inferior turbinate hypertrophy, excluding patients with refractory allergic/vasomotor/hypertrophic rhinitis. Turbinectomy and submucosal resection had elevated rates of crusting and epistaxis. Conservative treatments such as cryotherapy and submucous diathermy did not provide long-term results. Authors judged submucosal resection and radiofrequency ablation to have the most positive outcomes (decreased nasal resistance and preserved mucosal function (Sinno, 2016)).

The Sinno review did not distinguish results for children and adults separately. Other large reviews exclude pediatric patients (Acevedo, 2015; Mirza, 2020; Pedersen, 2018; Wright, 2020).

A large review included children younger than age 18 who underwent tonsillectomy/adenoidectomy (n = 72,043), turbinectomy (n = 639), or both (n = 3,079). Adding turbinectomy did not significantly raise rates of 14-day relevant revisits (9.4% versus 8.6%, $P = .11$) or hemorrhage requiring cauterization (1.5% versus 1.4%; $P = .64$) (Yeun, 2017).

A review of 3,669 patients younger than 18 years old with endoscopic sinus surgery showed turbinate reduction had no significant impact on rates of readmission within 30 days (McKeon, 2019).

In 2024, we removed threenon-seminal studies that were published more than 10 years ago. In addition, we found three studies that examined the efficacy, safety, and long-term outcomes of pediatric turbinate reduction surgery. A meta-analysis (n = 510) showed improved nasal patency post-surgery with similar success in various surgical techniques, highlighting the effectiveness of rhinoplasty in treating a nasal obstruction in the midterm (Alves de Sousa, 2023). A separate meta-analysis found that (n = 1411) durable symptom relief and low complications from turbinate surgery in long-term management of allergic rhinitis. Finally, a systematic review of 23 studies (n = 5206) found turbinate reduction generally safe and effective, particularly for cases resistant to medical treatment, with low complication rates such as crust formation (Aljerais, 2023).

References

On October 14, 2022, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were “nasal congestion,” “rhinitis,” “rhinosinusitis,” “turbinate hypertrophy,” “turbinectomy,” and “turbinoplasty.” We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

Abdullah B, Singh S. Surgical interventions for inferior turbinate hypertrophy: A comprehensive review of current techniques and technologies. *Int J Environ Res Public Health*. 2021;18(7):3441. Doi: 10.3390/ijerph18073441.

Acevedo JL, Camacho M, Brietzke SE. Radiofrequency ablation turbinoplasty versus microdebrider-assisted turbinoplasty: A systematic review and meta-analysis. *Otolaryngol Head Neck Surg*. 2015;153(6):951-956. Doi: 10.1177/0194599815607211.

Aljerais T. Efficacy and safety of turbinate reduction surgery in children: a systematic review. *Eur Arch Otorhinolaryngol*. Published online November 9, 2023. Doi:10.1007/s00405-023-08327-9

Alves de Sousa F, Santos M, Casanova M, et al. Pediatric inferior turbinate surgery: A review and meta-analysis of midterm nasal patency. *Int J Pediatr Otorhinolaryngol*. 2023;172:111661. Doi:10.1016/j.ijporl.2023.111661

Arganbright JM, Jenses EL, Mattingly J, Gap D, Chan KH. Utility of inferior turbinoplasty for the treatment of nasal obstruction in children: A 10-year review. *JAMA Otolaryngol Head Neck Surg.* 2015;141(10):901-904. Doi: 10.1001/jamaoto.2015.1560.

Calvo-Henriquez C, Capasso R, Martinez-Capoccioni G, et al. Safeness, subjective and objective changes after turbinate surgery in pediatric patients: A systematic review. *Int J Pediatr Otorhinolaryngol.* 2020;135:110128. Doi: 10.1016/j.ijporl.2020.110128.

Calvo-Henriquez C, Lechien JR, Mendez-Benegassi I, et al. Pediatric turbinate radiofrequency improves quality of life and rhinomanometric values. A prospective study. *Int J Pediatr Otorhinolaryngol.* 2022;154:111050. Doi: 10.1016/j.ijporl.2022.111050.

Ciprandi G, Tosca MA, Gallo F, Bassali GC, Ameli F. Turbinate hypertrophy in children with allergic rhinitis: Clinical relevance. *Acta Biomed.* 2020;91(1-S):43-47. Doi: 10.23750/abm.v91i1-S.9254.

Gary CC. Pediatric nasal surgery: Timing and technique. *Curr Opin Otolaryngol Head Neck Surg.* 2017;25(4):286-290. Doi: 10.1097/MOO.0000000000000378.

Isaacson G. Surgical treatment of pediatric rhinosinusitis. *Minerva Pediatr.* 2015;67(4):357-e368. <https://pubmed.ncbi.nlm.nih.gov/25369207/>.

Komshian SR, Cohen MB, Brook C, Levi JR. Inferior turbinate hypertrophy: A review of the evolution of management in children. *Am J Rhinol Allergy.* 2019;33(2):212-219. Doi: 10.1177/1945892418815351.

Maniaci A, Calvo-Henriquez C, Cammaroto G, et al. Pediatric inferior turbinate hypertrophy: diagnosis and management. A YO-IFOS consensus statement. *Laryngoscope.* 2024. Doi:10.1002/lary.30907

McKeon M, Medina G, Kawai K, Cunningham M, Adil E. Readmissions following ambulatory pediatric endoscopic sinus surgery. *Laryngoscope.* 2019;129(12):2681-2686. Doi: 10.1002/lary.27898.

Mirza AA, Alandejani TA, Shawli HY, Alsamel MS, Albakrei MO, Abdulazeem HM. Outcomes of microdebrider-assisted versus radiofrequency-assisted inferior turbinate reduction surgery: A systematic review and meta-analysis of interventional randomised studies. *Rhinology.* 2020;58(6):530-537. Doi: 10.4193/Rhin19.350.

National Institute for Health and Care Excellence. Radiofrequency tissue for turbinate hypertrophy. Interventional procedures guidance IPG495. <https://www.nice.org.uk/guidance/ipg495/chapter/2-Indications-and-current-treatments>. Published June 25, 2014.

Pedersen L, Schiöler L, Holmberg K, Ahlström Emanuelsson C, Hellgren J. Age and unplanned postoperative visits predict outcome after septoplasty: a national Swedish register study. *Int J Otolaryngol.* 2018;2(2018):2379536. Doi: 10.1155/2018/2379536.

Scheithauer MO. Surgery of the turbinates and "empty nose syndrome". *GMS Curr Top Otorhinolaryngol Head Neck Surg.* 2010;9:Doc03. Doi: 10.3205/cto000067.

Sedaghat AR. Chronic rhinosinusitis. *Am Fam Physician.* 2017;96(8):500-506. <https://pubmed.ncbi.nlm.nih.gov/29094889/>.

Seidman MD, Gurgel RK, Lin SY, et al. Clinical practice guideline: Allergic rhinitis. *Otolaryngol Head Neck Surg.* 2015;152(1 Suppl):S1-43. Doi: 10.1177/0194599814561600.

Sinno S, Mehta K, Lee Z-H, Kidwai S, Saadeh PB, Lee MR. Inferior turbinate hypertrophy in rhinoplasty: Systematic review of surgical techniques. *Plast Reconstr Surg*. 2016;138(3):419e-429e. Doi: 10.1097/PRS.0000000000002433.

Wald ER, Applegate KE, Bordley C, et al. Clinical practice guideline for the diagnosis and management of acute bacterial sinusitis in children aged 1 to 18 years. *Pediatrics*. 2013;132(1):e262-e280. Doi: 10.1542/peds.2013-1071.

Wright L, Grunzweig KA, Totonchi A. Nasal obstruction and rhinoplasty: A focused literature review. *Aesthetic Plast Surg*. 2020;44(5):1658-1669. Doi: 10.1007/s00266-020-01710-5.

Yuen SN, Leung PP, Funamura J, Kawai K, Roberson DW, Adil FA. Complications of turbinate reduction surgery in combination with tonsillectomy in pediatric patients. *Laryngoscope*. 2017;127(8):1920-1923. Doi: 10.1002/lary.26421.

Policy updates

1/2022: initial review date and clinical policy effective date: 2/2022

4/2023: Policy modified from investigational to medically necessary, references updated.

3/2024: Policy references updated.